

Noaedoxine

(Meclizine HCl, B6)

Tablets

INDICATIONS :

Management of nausea, vomiting and dizziness alone or associated with motion sickness and of nausea and vomiting due to pregnancy.

DOSAGE:

For Adults:

The recommended dose in 25 to 100mg Meclizine HCl daily, in divided dosage, depending upon clinical response . For the treatment of motion sickness the initial recommended dose of 25 to 50mg Meclizine HCl should be taken one hour prior to embarkation for protection against motion sickness. Thereafter, the dose may be repeated every 24 hours for the duration of the journey.

For children:

The recommended dosage are:

Upto 2 years : 5 to 10mg Meclizine HCl daily
2-5 years : 10 to 20mg Meclizine HCl daily
5-10 years : 15 to 30mg Meclizine HCl daily

خوراک:

برائے بالغان:

۲۵ تا ۱۰۰ ملی گرام روزانہ

بچوں کے لئے:

۲ سال تک : ۲۵ تا ۱۰ ملی گرام روزانہ

۲ سے ۵ سال تک : ۱۰ تا ۲۰ ملی گرام روزانہ

۵ سے ۱۰ سال تک : ۱۵ سے ۳۰ ملی گرام روزانہ

۲ سال سے کم عمر بچوں کے لئے نوایدوکسن ڈاکٹر کی

ہدایات کے بغیر استعمال نہ کریں۔

Noaedoxine in children under 2 year of age should be used only following the prescription of a physician.

CONTRAINDICATIONS :

Meclizine HCl is contraindicated in individuals who have shown a previous hypersensitivity to it.

SIDE EFFECTS:

Drowsiness , dry mouth and on rare occasion , blurred vision have been reported.

PRECAUTIONS:

Since drowsiness may , on occasion, occur with use of this drug, patients should be warned of this possibility and cautioned against driving vehicle or operating dangerous machinery.

Patient should avoid alcoholic beverages while taking the drug. Due to its potential anti-cholinergic action , this drug should be used with caution in patients with asthma , glaucoma , or enlargement of the prostate gland.

PRESENTATION:

Pack of 10x10 tablets, each tablet containing Meclizine HCl 25mg and Vitamin B6 50mg.

STORAGE CONDITION:

Do not store above 30°C. Protect from light. moisture & heat.

ہدایات: 30°C سے زیادہ درجہ حرارت پر نہ رکھیں۔

روشنی اور نمی سے بچائیں۔



Manufactured by :
SIGMA PHARMA
International (Pvt.) Ltd.
www.sigmapharma.com.pk

E-50, North Western
Industrial Zone,
Port Qasim Authority,
Karachi, Pakistan.

تمام دوائیں بچوں کی پہنچ سے دور رکھیں۔
دوا کو روشنی سے محفوظ رکھیں اور خشک جگہ پر رکھیں۔
ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔