## BLESSEL\_100mg/5ml. Injection IV



CINICAL PARAMACIONO'
Mechanism of Action
Following Intraversus administration, BLESEL (iron Sucree) is dissociated into iron and sucree and the iron is transported as a complex with
resident to large des including enythmod precupar cells. The iron in the precupar cells is incorporated into hemogoloin as the cells nature
Pharmaciónnetics
Inhabity adults streated with intravenous doors of BLESEL:
- Elimitation habilité to & Chous
Inhabity adults streated with intravenous doors of BLESEL:
- Elimitation habilité to & Chous
Inhabity adults streated with intravenous doors of BLESEL:
- Streated years apparent volume of distribution: 10.
- Steedy state apparent volume of distribution: 29.
Inhabity adults of the inhabity of the inhab

- Performed displant descreted channels idently disease (PDD-EXI) production sections are symptoposition.

DOSAGE AND ADMINISTRATION:
Dosage:
Administration of the production of the production of the state of the

Body Weight (kg)	Total number of BLESSEL (Iron Sucrose) ampoules to be administered (1 ampoule of BLESSEL (Iron Sucrose) corresponds to 5mL)			
	Hb 60g/L	Hb 75g/L	Hb 90g/L	Hb 105g/L
5	1.5	1.5	1.5	1
10	3	3	2.5	2
15	5	4.5	3.5	3
20	6.5	5.5	5	4
25	8	7	6	5.5
30	9.5	8.5	7.5	6.5
35	12.5	11.5	10	9
40	13.5	12	11	9.5
45	15	13	11.5	10
50	16	14	12	10.5
55	17	15	13	11
60	18	16	13.5	11.5
65	19	16.5	14.5	12
70	20	17.5	15	12.5
75	21	18.5	16	13
80	22.5	19.5	16.5	13.5
85	23.5	20.5	17	14
90	24.5	21.5	18	14.5

Children

Immined data for children under study, if there is a clinical need, it is recommended not to exceed 0.15mL BLESSEL (for Clinic limited data for children under study, if there is a clinical need, it is recommended not to exceed 0.15mL BLESSEL (for Clinic limited study or to three times per week depending on the hemoglobin level.

Maximum Date:

Introduction influence

Introduction influence

Introduction influence

The maximum allowed single dose is 7mg iron per kg body weight given once per week, but not exceeding 500mg iron.

The maximum allowed single dose is 20mg iron.

The maximum allowed single dose is 20mg iron.

The maximum allowed single dose is 200mg irron.

NDD-CED

ADD CED

* Dilution of BLESSEL		
Strength	0.9% w/v Sodium Chloride solution (Diluting Agent)	
BLESSEL 5mL (100mg)	100mL	
BLESSEL 10mL (200mg)	200mL	
Infusion Rate	·	
Dosage	Time	
BLESSEL 5mL (100mg)	15 minutes	
BLESSEL 10mL (200mg)	30 minutes	
BLESSEL 15mL (300mg)	1.5 hours	
BLESSEL 25mL (400mg)	2.5 hours	
**BLESSEL25mL (500mg)	3 hours	
***BLESSEL25mL (7mg iron/kg)	3.5 hours	
Its above 70kg, strict of 70kg and below. ous Injection: ous injection administered at a rate of 1mL undiluted solutic is (200mg iron) per injection. dministering a slow intravenous injection, a test dose of 1ml revise events cour within 1S minutes of completing the test	(20mg of iron) should be injected slowly over a period of 1 to	
nts above 70%; ents of 70%; and below. nous injection and below. nous injection administered at a rate of 1mt. undiluted solution set 2000mg (not) per injection. the set of the set of the set of the set of 1mt. where set of the set of 1mt. Life Tour Surceol is per administered during a hemolalysis : nors as those outlined for intravenous injection.	. (20mg of iron) should be injected slowly over a period of 1 to dose, then the remaining portion of the injection may be given	
Int above Yug,  star Stown Yug,  star O'Tilg, and bellow.  sto pictors administered at a rate of I'm, undiluted solution  solution injection, injection, injection, injection, and injection and injection and injection and injection, a set does of I'm, interess events occur with 15 minutes of controllegate the test  into dialyse:  (I'm o'successed jump be administered during a hemodalysis:  AMBOLATION of the first interess injection,  AMBOLATION outcomes of injection, or injection, and injection and injection.	. (20mg of iron) should be injected slowly over a period of 1 to dose, then the remaining portion of the injection may be give	
ins above Ying,  when yield a below,  sour hylectron  sour hylectron  sour hylectron  sour hylectron  sour hylectron  contribution  definition of the contribution  definition of the contribution  definition of the contribution of the contribution  definition of the contribution  when events occur with 15 minutes of completing the test  into dialyses.  res as those outlined for intravenous injectron,  when the contribution of the contribution  when the contribution of the contribution  is not attributable to into deficiency.  so not attributable to into indeficiency.	. (20mg of iron) should be injected slowly over a period of 1 to dose, then the remaining portion of the injection may be given	
Int above Ying,  see The Time In Time	. (20mg of iron) should be injected slowly over a period of 1 to dose, then the remaining portion of the injection may be give	
is above 70/Eq.  If of 70/Eq and below.  If of 70/Eq and 1/Eq.  If of 70/Eq.  If of	. (20mg of iron) should be injected slowly over a period of 1 to dose, then the remaining portion of the injection may be given	
is above 70/g, and is a dove 70/g, and below. It is a dove 70/g, and below. It is 70/g, and 10/g, and 10	Doing of Inco) should be injected slowly over a period of 1 to lose, then the remaining portion of the injection may be given session directly into the venous limb of the diabyser under the directly into the venous limb of the diabyser under the	
les (200mg unol per injection.  diministering a slow untreavours injection, a test dose of Imiteria events occur within 1.5 ministers of competiting the test inverse events occur within 1.5 ministers of competiting the test users as those outlined for intravenous injection.  June 3 and 10	n per minute (i.e., 5 minutes/ampoule) and not exceeding <b>BLI</b> (Jömg of iron) should be lejected diswly over a period of 1 to lose, then the remaining portion of the electron may be given session directly into the venous limb of the dialyser under the interval of the control of the dialyser under the hydrocol of reactions, which may be potentially fatal. Therefore,	

Page 1974 in a restrict.

Page LATIONS:

Page 1974 administered iron preparations can cause allergic or anaphylactoid reactions, which may be potentially fatal. Therefore, treatment of mild and serious allegar creations must be available.

Page 1974 iron administration should be avoided in parieties with hepatic dysfunction where iron overload is a precipitating factor. Careful monitoring of iron status is recommended to wood iron overload.

Page 1974 iron must be used with causion in case of acute or chronic infection.

Page 1974 iron process or page 1974 iron page 197

Thouse the second section of the second section of the second section of the sect

OVERTICALE:
Overtrapps should be treated with supportive measures and, if required, with an iron chelating agent.
INSTRUCTIONS:
Shows below 25C.
-Putter firm light & lout. Do not freeze.
-Putter fir

خوراک: ڈاکٹر کی ہدایت کےمطابق استعال کریں۔ بدایات: دواکو ۲۵ و کری سنتی کریدے کم درجه حرارت پر دھیں۔ روشی، گری اور مجمد مونے سے بچاکس ۔ بچوں کی کافی سے دورر کیس۔ انتاه: صرف دجيرٌ ذُوْاكمُّرُ كَ لَيْغِ كِمِطالِق فروشت كريں۔

