

68mm

68mm

145mm

145mm

SIGMADON

[Propyphenazone / Caffeine]

175mg/25mg Tablets

سگماڈون

۵۷۵ ملی گرام / ۲۵ ملی گرام کوبیاں

DESCRIPTION:

SIGMADON is effective and well tolerated analgesic and antipyretic which does not impair working capacity.

QUALITATIVE AND QUANTITATIVE COMPOSITION:

SIGMADON (Propyphenazone / Caffeine) is available for oral administration as:

SIGMADON 175mg/25mg Tablet
Each sugar coated tablet contains:
Propyphenazone BP175mg
Caffeine Anhydrous BP25mg
(Product Specs.: Sigma)

PHARMACOKINETICS:**Propyphenazone:**

The maximal plasma concentration is reached within 1 to 3 hours. The elimination half life is 1 to 5 hours. About 0.6% of the unchanged drug is excreted in the urine.

Caffeine:

Caffeine is absorbed readily after oral administration. Maximal plasma concentrations are achieved within one hour and the plasma half-life is about 3.5 hours. 65 - 80% of administered caffeine is excreted in the urine as 1-methyluric acid and 1-methylxanthine.

INDICATIONS:

SIGMADON is indicated for, the following:

- Relieves Fever
- Pain of Various Origins:
 - Headache
 - Toothache
 - Dysmenorrhoea
 - Joint Pain
 - Muscle Pain
 - Backache

DOSAGE AND ADMINISTRATION:

The coated tablets should be swallowed with some liquid without chewing.

Adults:

1 or 2 Tablets at a time, if necessary this dose may be repeated over the day once or twice.

Maximum daily dose 4 Tablets.

Children (from 6 years of age):

1 Tablet at a time.

Maximum daily dose 1-2 Tablets.

CONTRAINDICATION:

Known hypersensitivity to pyrazolone derivatives.

PRECAUTIONS:

- **SIGMADON** should be reserved for the treatment of painful episodes. It is not intended for prolonged treatment.
- Caution is required in cases of severe kidney or liver disorders.

PREGNANCY AND LACTATION:

SIGMADON should be used during pregnancy or lactation only on advice of the doctor.

ADVERSE REACTIONS:

- Occasionally dizziness and nausea may occur.
- Very rarely hypersensitivity reactions including skin rashes and anaphylaxis-like symptoms have been observed.
- With products containing pyrazolone derivatives, blood dyscrasias have been reported in isolated cases.

OVERDOSAGE:

- Symptoms: Abdominal pain, nausea, vomiting, somnolence, excitation, convulsions, tachycardia and coma.
- Treatment: Treatment is symptomatic; gastric lavage, if indicated, should be performed.

INSTRUCTIONS:

- Store below 30°C.
- Protect from light & moisture.
- Keep out of reach of children.

To be sold on prescription of a registered medical practitioner only.

HOW SUPPLIED:

SIGMADON (Propyphenazone / Caffeine) 175mg/25mg Tablets are available in pack of 20x10's.

نوراک: ٹیکس کی یہ ایت کے مطابق استعمال کریں۔
ہدایات: دوا کو ۳۰ ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔
روشنی اور نمی سے بچائیں۔ پھول کی پتھری سے دور رکھیں۔
صرف رجسٹرڈ ڈاکٹر کے نسخے کے مطابق فروخت کریں۔

**SIGMA
PHARMA**

Manufactured by:

**SIGMA PHARMA
International (Pvt.) Ltd.**

www.sigmapharma.com.pk

E-50, North Western
Industrial Zone,
Port Qasim Authority,
Karachi, Pakistan.